

Quantum mechanics on Riemannian Manifold in Schwinger's Quantization Approach III

Chepilko Nicolai Mikhailovich

Physics Institute of the Ukrainian Academy of Sciences, Kyiv-03 028, Ukraine

e-mail: chepilko@zeos.net

Romanenko Alexander Victorovich

Kyiv Taras Shevchenko University, Department of Physics, Kyiv-03 022, Ukraine

e-mail: ar@ups.kiev.ua

February 1, 2008

Abstract

Using extended Schwinger's quantization approach quantum mechanics on a Riemannian manifold M with a given action of an intransitive group of isometries is developed. It was shown that quantum mechanics can be determined unequivocally only on submanifolds of M where G acts simply transitively (orbits of G -action). The remaining part of degrees of freedom can be described unequivocally after introducing some additional assumptions. Being logically unmotivated, these assumptions are similar to canonical quantization postulates. Besides this ambiguity that has a geometrical nature there is undetermined gauge field of $\hbar$ (or higher) order, vanishing in the classical limit $\hbar \rightarrow 0$.

1 Introduction

The purpose of the present paper is to continue a series of our works devoted to a generalization of Schwinger's quantization approach for the case of a Riemannian manifold with a group structure (determined by its metric).

In this paper we turn to the most problematic case (in physical meaning) of the formulation of quantum mechanics on a Riemannian manifold M with the intransitive group of isometries G . Partially our consideration is based on our previous results [2], [3]. But, in contradiction to them, the main feature of the present paper is the fact that the dimension of the manifold M is bigger than the dimension of the group of isometries G . This leads to the decomposition of degrees of freedom describing a point particle on M into two sets, related to the group G and to the quotient space M/G . The last manifold has not a global group structure in a general case. Due to this reason it happens that the quantum mechanics is completely defined only on submanifolds of M , which are isomorphic to G (the orbits of action of G on M). The dynamical equations for the other degrees of freedom, connected with M/G , can be unequivocally determined after introducing some new assumptions expanding Schwinger's scheme from outside, that are similar to canonical quantization postulates and can not be motivated logically. Nevertheless, having introduces them, we find that in quantum mechanics

there is an abelian gauge field of $\hbar$ (or higher) order, that remains undetermined and vanishes in the classical limit $\hbar \rightarrow 0$.

2 Structure of Manifold with Non-transitive Group of Isometries

Let us consider p -dimensional Riemannian manifold M equipped with the metric $\{\eta_{MN} : M, N = \overline{1, p}\}$ in which the action of n -dimensional intransitive group of isometries G is given ($n < p$). This means, according to [1], that the Killing equations

$$v^P \partial_P \eta_{MN} + \eta_{MP} \partial_N v^P + \eta_{PN} \partial_M v^P = 0 \quad (2.1)$$

have n independent solutions $\{v_A^N : N = \overline{1, p}; A = \overline{1, n}\}$.

The set of vector fields $\{v_A^M \partial_M : A = \overline{1, n}\}$ describes the representation of the Lie algebra $\text{Lie}(G)$ of the Lie Group G acting on M . Due to this fact each the vector from this set obeys the equation

$$v_A^N \partial_N v_B^M - v_B^N \partial_N v_A^M = C^C{}_{AB} v_C^M, \quad (2.2)$$

where $C^C{}_{AB}$ are the structure constants of G . This equality provides the following equation to have $m = p - n$ independent solutions

$$v_A^N \partial_N \varphi(x) = 0. \quad (2.3)$$

Let us consider a new coordinate system

$$\bar{x}^N := (\varphi^\alpha(x), x^\mu) \quad (2.4)$$

$$N = \overline{1, p}, \quad \alpha = \overline{1, m}, \quad \mu = \overline{m+1, p}$$

where $\{\varphi^\alpha(x) : \alpha = \overline{1, m}\}$ denote independent solutions of (2.3). In these coordinates the Killing vectors take the form:

$$\bar{v}_A^N = (v_A^M \partial_N \varphi^\alpha, v_A^\mu) \equiv (0, v_A^\mu). \quad (2.5)$$

Further we will assume that such a coordinate system denoted by $\{x^N\}$ on M is given and all the geometrical objects are expressed in terms of it. The Killing vector can be rewritten as

$$v_i^M = \delta_\mu^M v_i^\mu, \quad (2.6)$$

where the group index $i = \overline{m+1, p}$ is used instead of $A = \overline{1, n}$ (the indices $i, j, k, \dots$ play the same role as $A, B, C, \dots$).

We introduce the inverse of the matrix $\{v_i^\mu : \mu = \overline{m+1, p}, i = \overline{m+1, p}\}$ as

$$e_\mu^i v_j^\mu = \delta_j^i, \quad e_\mu^i v_i^\nu = \delta_\mu^\nu. \quad (2.7)$$

Using (2.2), (2.6), (2.7) one can prove that the matrix $\{e_\mu^i\}$ obeys the Maurer-Cartan equation

$$\partial_\mu e_\nu^i - \partial_\nu e_\mu^i = -C^i{}_{jk} e_\mu^j e_\nu^k. \quad (2.8)$$

It is easy to obtain the following equations by differentiating

$$\begin{aligned}
D_M v_i^\mu &= \partial_M v_i^\mu + \Lambda_{M\nu}^\mu v_i^\nu = 0 \\
D_M e_\mu^i &= \partial_M e_\mu^i - \Lambda_{M\mu}^\nu e_\nu^i = 0
\end{aligned} \tag{2.9}$$

where the object

$$\Lambda_{M\nu}^\mu = v_i^\mu \partial_M e_\nu^i \tag{2.10}$$

corresponds to the right group connection on M (see [1]).

Using (2.7), (2.8) one can easily prove the following property of $\Lambda_{M\nu}^\mu$:

$$\Lambda_{\mu\nu}^\sigma - \Lambda_{\nu\mu}^\sigma = -C_{jk}^i v_i^\sigma e_\mu^j e_\nu^k. \tag{2.11}$$

Using the relations obtained above we can write down the functional features of the metric in the special coordinate system, described by (2.4). To do it we transform the Killing equation (2.1) into new coordinate description:

$$\begin{aligned}
\partial_\mu \eta^{\alpha\beta} &= 0, \\
\partial_\mu \eta_{\alpha\nu} &= \eta_{\alpha\sigma} \Lambda_{\nu\mu}^\sigma - \eta_{\nu\sigma} \Lambda_{\alpha\mu}^\sigma, \\
\partial_\mu \eta_{\alpha\beta} &= \eta_{\alpha\nu} \Lambda_{\beta\mu}^\nu + \eta_{\beta\nu} \Lambda_{\alpha\mu}^\nu.
\end{aligned} \tag{2.12}$$

The matrix $\{\eta_{\mu\nu} : \mu, \nu = \overline{m+1, p}\}$ have the sense of the metric on the orbit of the action of G on M , that is isomorphic to G .

Making an assumption that $\{\eta_{\mu\nu}\}$ is non-degenerate matrix we introduce the following objects:

$$A_\alpha^\mu = g^{\mu\nu} \eta_{\alpha\nu}, \quad g^{\mu\nu} \eta_{\nu\sigma} = \delta_\sigma^\mu. \tag{2.13}$$

Then $\eta_{\alpha\mu} = g_{\mu\nu} A_\alpha^\nu$ and

$$D_\mu A_\alpha^\nu = \partial_\mu A_\alpha^\nu + \Lambda_{\mu\sigma}^\nu A_\alpha^\sigma = \Lambda_{\alpha\mu}^\nu, \tag{2.14}$$

as it follows from (2.12).

Due to (2.13), (2.14) the second equation in (2.12) is equivalent to

$$\partial_\mu (\eta_{\alpha\beta} - g_{\mu\nu} A_\alpha^\mu A_\beta^\nu) = 0. \tag{2.15}$$

Hence, the matrix

$$g_{\alpha\beta} = \eta_{\alpha\beta} - g_{\mu\nu} A_\alpha^\mu A_\beta^\nu \tag{2.16}$$

depends only on coordinates $\{x^\alpha : \alpha = \overline{1, m}\}$ and is independent on $\{x^\mu : \mu = \overline{m+1, p}\}$. The metric tensor $\{\eta_{MN}\}$, describing the manifold M can be rewritten in terms of objects $g_{\mu\nu}$, A_α^μ , $g_{\alpha\beta}$ as follows

$$\{\eta_{MN}\} = \begin{pmatrix} g_{\alpha\beta} + g_{\rho\sigma} A_\alpha^\rho A_\beta^\sigma & g_{\mu\rho} A_\beta^\rho \\ A_\alpha^\rho g_{\rho\nu} & g_{\mu\nu} \end{pmatrix}, \tag{2.17}$$

$$\{\eta^{MN}\} = \begin{pmatrix} g^{\alpha\beta} & -A_\gamma^\mu g^{\gamma\beta} \\ -g^{\alpha\gamma} A_\beta^\nu & g^{\mu\nu} + g^{\gamma\delta} A_\gamma^\mu A_\delta^\nu \end{pmatrix}. \tag{2.18}$$

This decomposition is similar to Kaluza-Klein one. In (2.18) $\{g^{\alpha\beta}\}$ denotes the inverse of $\{g_{\alpha\beta}\}$. If $\{\eta_{MN}\}$ and $\{g_{\mu\nu}\}$ are non-degenerate matrices the matrix $\{g^{\alpha\beta}\}$ exists due to the following property

$$\det \{\eta_{MN}\} = \det \{g_{\alpha\beta}\} \cdot \det \{g_{\mu\nu}\}$$

The functional properties of the metric (2.17) are determined by the following equations

$$\begin{cases} \partial_\mu g_{\alpha\beta} &= 0 \\ \partial_\rho g_{\mu\nu} &= g_{\mu\sigma} \Lambda_{\rho\mu}^\sigma + g_{\nu\sigma} \Lambda_{\rho\nu}^\sigma \\ \partial_\mu A_\alpha^\nu &= \Lambda_{\alpha\mu}^\nu - A_\alpha^\rho \Lambda_{\rho\mu}^\nu \end{cases} \quad (2.19)$$

By its written form, the matrices (2.17), (2.18) correspond to analogous ones in [3], but their meaning is different.

From the geometrical point of view the construction we are considering corresponds to the principal bundle with a total space M , a base space M/G and a structure group G . The coordinates $\{x^\alpha\}$ from the set $\{x^M\} = \{x^\alpha, x^\mu\}$ describes the local coordinate system in M/G , the matrix $\{g_{\alpha\beta}\}$ — the metric tensor of M/G .

The projection map of the principal bundle $(M/G) (M, G)$ has the form

$$\begin{aligned} p: M &\longrightarrow M/G \\ \{x^\alpha, x^\mu\} &\longrightarrow \{x^\alpha\}. \end{aligned}$$

The vector fields

$$\hat{D}_i = v_i^\mu \partial_\mu, \quad \hat{D}_\alpha = \partial_\alpha - A_\alpha^\mu \partial_\mu \quad (2.20)$$

form the bases of vertical and horizontal fields correspondingly. One can observe it from

$$dp(\hat{D}_i|_{\{x^\alpha, x^\mu\}}) = 0, \quad dp(\hat{D}_\alpha|_{\{x^\alpha, x^\mu\}}) = \partial_\alpha|_{\{x^\alpha\}}. \quad (2.21)$$

The objects $\{A_\alpha^\mu\}$ defined above correspond to the Lie (G) -valued connection 1-form on M :

$$\omega^i = e_\mu^i (dx^\mu + A_\alpha^\mu dx^\alpha). \quad (2.22)$$

Now we write down some useful relations for the Riemannian geometry on M , which describe its features explicitly.

Let ∇_α be a standard covariant derivative constructed with the metric $\{g_{\alpha\beta}\}$. Define the operator on M/G that generalize the operator $\hat{\nabla}_\alpha$ as

$$\hat{\nabla}_\alpha B_\beta = \hat{D}_\alpha B_\beta - \Gamma_{\alpha\beta}^\gamma B_\gamma \quad (2.23)$$

for some vector field B_α on M/G depending on coordinates $\{x^\mu\}$. Then

$$[\hat{\nabla}_\alpha, \hat{\nabla}_\beta] B_\gamma = [\hat{D}_\alpha, \hat{D}_\beta] B_\gamma - R_{\alpha\beta\gamma}^\delta B_\delta, \quad (2.24)$$

where $R_{\alpha\beta\gamma}^\delta$ is the curvature tensor for the metric $g_{\alpha\beta}$.

The explicit expression for the commutator $[\hat{D}_\alpha, \hat{D}_\beta]$ can be obtained from its action on a scalar function $f(x^\alpha, x^\mu)$. Performing the simple calculation we can observe that

$$[\hat{D}_\alpha, \hat{D}_\beta] f = -(\partial_\alpha A_\beta^\mu - \partial_\beta A_\alpha^\mu) \partial_\mu f + (A_\alpha^\mu \partial_\mu A_\beta^\nu - A_\beta^\mu \partial_\mu A_\alpha^\nu) \partial_\nu f. \quad (2.25)$$

Using (2.19) in (2.25):

$$A_\alpha^\mu \partial_\mu A_\beta^\nu - A_\beta^\mu \partial_\mu A_\alpha^\nu = -C_{ik}^i A_\alpha^j A_\beta^k v_i^\nu + (\partial_\alpha A_\beta^\nu - \partial_\beta A_\alpha^\nu) + (\partial_\alpha A_\beta^i - \partial_\beta A_\alpha^i) v_i^\nu, \quad (2.26)$$

where

$$A_\alpha^i = e_\mu^i A_\alpha^\mu, \quad \partial_\mu A_\alpha^i = -C_{jk}^i e_\mu^j A_\alpha^k + \partial_\alpha e_\mu^i. \quad (2.27)$$

Due to (2.26), (2.27) the expression (2.25) can be rewritten as

$$[\hat{D}_\alpha, \hat{D}_\beta]f = -F_{\alpha\beta}^i \hat{D}_i f, \quad \hat{D}_i = v_i^\mu \partial_\mu. \quad (2.28)$$

$$F_{\alpha\beta}^i = \partial_\alpha A_\beta^i - \partial_\beta A_\alpha^i + C_{jk}^i A_\alpha^j A_\beta^k. \quad (2.29)$$

Observing (2.28)-(2.29) we can draw a conclusion that the objects $\{A_\alpha^i\}$ can be interpreted as gauge fields defined on M/G with the strength tensor $F_{\alpha\beta}^i$.

The other relations between the basic fields have the form

$$[\hat{D}_i, \hat{D}_j] = C_{ij}^k \hat{D}_k, \quad [\hat{D}_\alpha, \hat{D}_i] = 0.$$

At the end of this section we consider classes of coordinate transformations $x^\mu \rightarrow \bar{x}^\mu$ preserving the form of the metric (2.17)-(2.18) and its functional structure. Evidently, one such a class consists of coordinate transformations on the orbit of the action of G :

$$\begin{cases} \bar{x}^\alpha &= x^\alpha \\ \bar{x}^\mu &= \bar{x}^\mu(x^\nu) \end{cases} \quad (2.30)$$

Under such a transformation $g_{\mu\nu}$, A_α^μ behave as a tensor and a covariant vector correspondingly (the geometric properties are described by $\mu, \nu, \dots$). In the same time $g_{\alpha\beta}$ transforms as a scalar.

The coordinate change on M/G

$$\begin{cases} \bar{x}^\alpha &= \bar{x}^\alpha(x^\beta) \\ \bar{x}^\mu &= x^\mu \end{cases} \quad (2.31)$$

is of the same class.

The objects $g_{\alpha\beta}$, A_α^μ transform as a tensor and a contravariant vector correspondingly (the geometric properties are described by $\alpha, \beta, \dots$), while $g_{\alpha\beta}$ transforms as a scalar.

The third class of transformations is described by

$$\begin{cases} \bar{x}^\alpha &= x^\alpha \\ \bar{x}^\mu &= x^\mu + \varphi^\mu(x^\alpha) \end{cases} \quad (2.32)$$

The main geometrical objects transform under (2.32) as

$$\bar{g}_{\mu\nu}(\bar{x}) = g_{\mu\nu}(x), \quad \bar{g}_{\alpha\beta}(\bar{x}) = g_{\alpha\beta}(x), \quad \bar{A}_\alpha^\mu(\bar{x}) = A_\alpha^\mu - \partial_\alpha \varphi^\mu(x^\alpha).$$

An arbitrary vector F_M on M can be decomposed into two objects that are invariant under the transformation (2.32):

$$\mathcal{F}_\alpha = F_\alpha - A_\alpha^\mu F_\mu, \quad \mathcal{F}_\mu = F_\mu.$$

This decomposition means the extraction of the horizontal part of F_M .

3 Lagrangian and Variational Principle

Constructing the quantum mechanics on a Riemannian manifold M with the non-transitive group of isometries G in terms of variational principle, as in our previous papers, we assume that the coordinate operators x^M form the total set of commuting observables.

The quantum Lagrangian can be written as

$$L = \frac{1}{2} \dot{x}^M \eta_{MN}(x) \dot{x}^N - U_q(x), \quad (3.33)$$

where $U_q(x)$ is some function that provides the scalar transformation law of L under a general non-degenerate coordinate transformation $x \rightarrow \bar{x} = \bar{x}(x)$ (see [2], [3]). As it has been pointed out in [2], its explicit expression can be determined in the case when the commutator $[x^M, \dot{x}^N]$ is the function of only $\{x^M\}$. This is a standard assumption in the formulation of the quantum mechanics on a Riemannian manifold (see [4] and motivation in [2]). The function U_q appears as the value of $\hbar^2$ -order.

In the special coordinate system, introduced in (2.4), the metric tensor $\{\eta_{MN}\}$ receives the form that is similar to one appearing in Kaluza-Klein theories. The Lagrangian can be written as

$$L = \frac{1}{2} (\dot{x}^\mu + \dot{x}^\alpha A_\alpha^\mu) g_{\mu\nu} (\dot{x}^\nu + A_\beta^\nu \dot{x}^\beta) + \frac{1}{2} \dot{x}^\alpha g_{\alpha\beta} \dot{x}^\beta - U_q(x). \quad (3.34)$$

The Euler-Lagrange dynamical equations, according to [2], can be derived from the action principle in the form of the variational equation $\delta L = 0$, that corresponds to the infinitesimal coordinate variation $x^M \rightarrow x^M + \delta x^M(x)$.

As it has been developed in [2], the equality $\delta L = 0$ holds if and only if the variations δx^M are Killing vectors. In the present case, in contradiction to one investigated in [2] and [3], the dimension of the manifold M is less than the number of linear independent Killing vectors $\{v_A^M : M = \overline{1, p}, A = \overline{1, n}\}$ (i.e. $n < p$).

Extracting the total time derivative, as in [2], we can rewrite the Lagrangian as

$$\delta L = \frac{d}{dt} (p_M \circ \delta x^M) - \dot{p}_M \circ \delta x^M + \frac{1}{2} \dot{x}^M (\delta x^P \partial_P \eta_{MN}) \dot{x}^N - \delta x^M \partial_M U_q + \frac{1}{2} \left[\delta x^M, \frac{d}{dt} [\dot{x}^N, \eta_{MN}] \right], \quad (3.35)$$

where $p_M = \eta_{MN} \circ \dot{x}^N$ means the momentum operator on M .

In accordance with [2], the object

$$G = p_M \circ \delta x^M \quad (3.36)$$

have the sense of the generator of permissible variations.

Further we decompose the variation δx^M in terms of the basis of linear independent Killing vectors (2.6):

$$\delta x^M = v_i^M \varepsilon^i, \quad \varepsilon^i = \text{const}, \quad (3.37)$$

This allow us to rewrite (3.36) as

$$G = (p_M \circ v_i^M) \varepsilon_i = p_i \varepsilon^i, \quad p_i := p_M \circ v_i^M. \quad (3.38)$$

In the special coordinate system (2.4) the objects $\{p_i\}$ have the form

$$p_i = p_\mu \circ v_i^\mu, \quad p_\mu = \eta_{\mu M} \circ \dot{x}^M = g_{\mu\nu} \circ (\dot{x}^\nu + A_\alpha^\nu \circ \dot{x}^\alpha). \quad (3.39)$$

4 Algebra of Commutation Relations

Following the procedure presented in [2], we derive the commutation relations for quantum theory on M performing investigation of the properties of permissible variations connected with action of the group G of isometries on M . From the definition of permissible variations in terms of the representation of G on M (in the special coordinate system) we have

$$\delta_i x^\mu = v_i^\mu = \frac{1}{i\hbar} [x^\mu, p_i] . \quad (4.40)$$

$$\delta_i x^\alpha = 0 = \frac{1}{i\hbar} [x^\alpha, p_i] . \quad (4.41)$$

(here $\delta x^\mu = \varepsilon^i \delta_i x^\mu$). Using the assumption about the commutativity of the coordinates $\{x^M\}$ and taking into account the fact that the matrix $\{v_i^\mu\}$ is non-degenerate (i.e. $\det v_i^\mu \neq 0$), we can conclude from (4.40), (4.41) that

$$[x^\mu, p_\nu] = i\hbar \delta_\nu^\mu , \quad [x^\alpha, p_\mu] = 0 . : \quad (4.42)$$

The variation of the arbitrary function f which depends on the coordinates $\{x^M\}$ can be defined as follows

$$\delta_i f(x) = v_i^\mu \partial_\mu f(x) = \frac{1}{i\hbar} [f, p_\mu] . \quad (4.43)$$

Hence

$$[f(x), p_\mu] = i\hbar \partial_\mu f(x) . \quad (4.44)$$

The variations of the velocity operators are

$$\delta_i \dot{x}^\mu = \frac{dv_i^\mu}{dt} = \dot{x}^M \circ \partial_M v_i^\mu = \frac{1}{i\hbar} [\dot{x}^\mu, p_i] , \quad (4.45)$$

$$\delta_i \dot{x}^\alpha = \frac{dv_i^\alpha}{dt} = 0 = \frac{1}{i\hbar} [\dot{x}^\alpha, p_i] . \quad (4.46)$$

Using (3.38), (3.39) we can express the velocity operators $\{\dot{x}^\mu\}$ in (4.45) in terms of the momentum operators $\{p_\mu\}$. Finally, we have

$$[p_\mu, p_i] = -i\hbar p_\nu \circ \partial_\mu v_i^\nu . \quad (4.47)$$

The commutator (4.47) can be presented in another form, taking into account (2.2)

$$[p_i, p_j] = -i\hbar C^i_{ik} p_k . \quad (4.48)$$

Similarly, due to (2.2) and (4.44), we can write

$$[p_\mu, p_\nu] = 0 . \quad (4.49)$$

The commutators (4.42), (4.44), (4.48) and (4.49) completely define quantum mechanics on the orbit of the action of G on M , that has been analyzed in [2].

Further, returning to the manifold M/G , we define the momentum operator

$$p_\alpha = \eta_{\alpha M} \circ \dot{x}^M \quad (4.50)$$

and the auxiliary operator

$$\pi_\alpha = g_{\alpha\beta} \circ \dot{x}^\beta = p_\alpha - A_\alpha^\mu \circ p_\mu. \quad (4.51)$$

Due to $\partial_\mu g_{\alpha\beta} = 0$, the objects p_μ and π_μ are connected by the following commutation relations

$$[p_\alpha, p_i] = -i\hbar \circ \partial_\alpha v_i^\mu, \quad [\pi_\alpha, p_i] = 0. \quad (4.52)$$

These are all the commutators which can be obtained immediately from the generator of permissible variations (3.38). Note, that presented above commutation relations are form-invariant under a general coordinate transformation $x^M \rightarrow \bar{x}^M = \bar{x}^M(x)$. The derivation of the remaining commutators requires the usage of operator equalities and introducing some additional assumptions.

It follows from the basic assumptions that the commutator between coordinate and momentum operators on M is the function of only $\{x^M\}$, i.e.

$$[x^M, p_N] = i\hbar B_N^M(x). \quad (4.53)$$

Due to the commutation relations obtained above the parts of the matrix B is already defined, namely $B_\nu^M = \delta_\nu^M$. Then we can write

$$B = \begin{pmatrix} B_\beta^\alpha & B_\beta^\mu \\ 0 & \delta_\nu^\mu \end{pmatrix} \quad (4.54)$$

where the unknown functions B_β^α , B_β^μ can be written in the following form motivated by the correspondence principle

$$B_\beta^\alpha = \delta_\beta^\alpha + b_\beta^\alpha, \quad B_\alpha^\mu = b_\alpha^\mu. \quad (4.55)$$

The new unknown objects b_β^α and b_α^μ in (4.54) are the functions of x of $\hbar^2$ (or higher) order. To derive their operator properties we can use the commutator of the structure equation

$$v_i^\mu \partial_\mu v_j^\nu - v_j^\mu \partial_\mu v_i^\nu = C^k_{ij} v_k^\nu \quad (4.56)$$

with the momentum operator p_α . Using the relation

$$[v_i^\mu, p_j] = i\hbar \hat{D}_j v_i^\mu \quad (4.57)$$

(the “long derivative” $\hat{D}_i$ was introduced in (2.20), (2.21)) we can rewrite (4.56) as

$$[v_i^\mu, p_j] - [v_j^\mu, p_i] = -C^k_{ij} v_k^\mu. \quad (4.58)$$

Therefore, using the Jacobi identity, we arrive to the following operator equality

$$[[v_i^\mu, p_j], p_\alpha] = -\hbar^2 \partial_\nu v_i^\mu \partial_\alpha v_j^\nu + i\hbar \hat{D}_j [v_i^\mu, p_\alpha]. \quad (4.59)$$

Further, taking the antisymmetrization of (4.59) with respect to the indices i, j we can find the equation

$$\hat{D}_i \varphi^\mu_{j\alpha} - \hat{D}_j \varphi^\mu_{i\alpha} = C^k_{ij} \varphi^\mu_{k\alpha} \quad (4.60)$$

where we introduce the unknown function of coordinates defined as

$$\varphi^\mu_{i\alpha} = \frac{1}{i\hbar} [v_i^\mu, p_\alpha] - \partial_\alpha v_i^\mu. \quad (4.61)$$

Taking into account the structure of the operator $\hat{D}_i$ we can conclude that the solution of (4.60) has the form

$$\varphi^\mu_{i\alpha} = \hat{D}_i \varphi^\mu_\alpha, \quad (4.62)$$

where φ^μ_α is a new unknown function of $\{x^M\}$.

Making the substitution $p_i \equiv v_i^\mu \circ p_\mu$ into (4.52) we obtain

$$[p_\alpha, p_\mu] = i\hbar \partial_\mu \varphi^\nu_\alpha \circ p_j. \quad (4.63)$$

Now we are going to show the relation between the objects φ^μ_α and B^ν_α . To do it let us write down the Jacobi identities including the coordinate operators x^M and momentum operators p_α , p_μ :

$$\frac{1}{i\hbar} [x^\alpha, [p_\beta, p_\mu]] \equiv 0 = \frac{1}{i\hbar} ([x^\alpha, p_\beta], p_\mu) - [[x^\alpha, p_\mu], p_\beta] = i\hbar \partial_\mu B^\alpha_\beta. \quad (4.64)$$

$$\frac{1}{i\hbar} [x^\mu, [p_\alpha, p_\nu]] \equiv i\hbar \partial_\nu \varphi^\mu_\alpha = \frac{1}{i\hbar} ([x^\mu, p_\alpha], p_\nu) - [[x^\mu, p_\nu], p_\alpha] = i\hbar \partial_\nu B^\mu_\alpha. \quad (4.65)$$

From (4.64) we can draw a conclusion about the independence of B^α_β (and, in consequence, b^α_β) on x^μ , because

$$\partial_\mu B^\alpha_\beta = \partial_\mu b^\alpha_\beta = 0. \quad (4.66)$$

At the same time, it follows from (4.65) that the differential connection between φ^μ_α and B^μ_α (and, in consequence, b^μ_α) is

$$\partial_\mu \varphi^\nu_\alpha = \partial_\mu B^\nu_\alpha = \partial_\mu b^\nu_\alpha. \quad (4.67)$$

The object φ^μ_α appears in all the formulae we deal with under the derivative operator ∂_μ . Hence, using (4.67) we can identify φ^μ_α with b^μ_α , i.e. take

$$\varphi^\mu_\alpha(x) = b^\mu_\alpha(x). \quad (4.68)$$

Due to the structure equation the commutator for v_i^μ and p_α reads

$$\frac{1}{i\hbar} [v_i^\mu, p_\alpha] = \partial_\alpha v_i^\mu + v_i^\nu \partial_\nu b^\mu_\alpha. \quad (4.69)$$

On the other hand, in accordance with (4.53), this commutator is equal to

$$\frac{1}{i\hbar} [v_i^\mu p_\alpha] = B^\mu_\alpha \partial_M v_i^\mu = \partial_\alpha v_i^\mu + b^\mu_\alpha \partial_M v_i^\mu. \quad (4.70)$$

By comparing the equations (4.69) and (4.70) we obtain the equation for the objects $\{b^M_\alpha\}$:

$$v_i^M \partial_M b^\mu_\alpha - b^\mu_\alpha v_i^M = 0, \quad \partial_\mu b^\alpha_\beta = 0. \quad (4.71)$$

This equation has the transparent geometrical meaning. In accordance with [1], the object $\{b^M_\alpha\}$ determines vector fields on M , that generate the one-parametric group of coordinate

transformations, commuting with isometries (isometric transformations). In a general case these transformations does not form the representation of an $p - n$ -dimensional group.

As the solutions of (4.71), the objects are not determined unequivocally. If b_α^M denotes the solution of (4.71), its linear combination

$$\bar{b}_\alpha^M(x) = \varepsilon_\alpha^\beta(x^\gamma) b_\beta^M(x) \quad (4.72)$$

also obeys the equation (4.71) for arbitrary functions $\varepsilon_\alpha^M = \varepsilon_\alpha^M(x^\gamma)$.

Therefore, the remaining commutation relations are

$$\begin{aligned} [x^\alpha, p_\beta] &= i\hbar (\delta_\beta^\alpha + b_\beta^\alpha), \\ [x^\mu, p_\beta] &= i\hbar b_\beta^\mu, \\ [p_\mu, p_\alpha] &= -i\hbar \partial_\mu b_\alpha^\nu \circ p_\nu, \end{aligned} \quad (4.73)$$

where b_α^M obeys the equation (4.71).

In order to construct the self-contained algebra of commutation relations for quantum mechanics on M we have to obtain the explicit form of $[p_\alpha, p_\beta]$. In a general case we can write

$$[p_\alpha, p_\beta] = i\hbar (F^M_{\alpha\beta} \circ p_M + \varphi_{\alpha\beta}), \quad (4.74)$$

where $F^M_{\alpha\beta}$, $\varphi_{\alpha\beta}$ are unknown tensors of coordinates on M . To determine these objects we use the Jacobi identities for operators x^α , p_β , p_γ and x^μ , p_α , p_β . After simple calculation we find that

$$F^\gamma_{\alpha\beta} B_\gamma^\delta = - (B_\alpha^\gamma \partial_\gamma B_\beta^\delta - B_\beta^\gamma \partial_\gamma B_\alpha^\delta), \quad (4.75)$$

$$F^\mu_{\alpha\beta} + B_\gamma^\mu F^\gamma_{\alpha\beta} = - (B_\alpha^M \partial_M B_\beta^\mu - B_\beta^M \partial_M B_\alpha^\mu). \quad (4.76)$$

These equations allow us to define $F^M_{\alpha\beta}$ as a function of B_α^M .

To investigate the functional properties of the object $\varphi_{\alpha\beta}(x)$ it is convenient to introduce the following operators

$$\bar{p}_\alpha = p_\alpha - b_\alpha^\mu \circ p_\mu. \quad (4.77)$$

According (4.74), the operators (4.77) satisfy the following commutation relations

$$[\bar{p}_\alpha, p_\mu] = 0, \quad [\bar{p}_\alpha, \bar{p}_\beta] = i\hbar (F^\gamma_{\alpha\beta} \circ \bar{p}_\gamma + \varphi_{\alpha\beta}). \quad (4.78)$$

From the relations

$$[p_\mu, [\bar{p}_\alpha, \bar{p}_\beta]] = 0, \quad \partial_\mu F^\gamma_{\alpha\beta} = 0, \quad (4.79)$$

obtained from (4.77), (4.78) we can conclude that

$$\partial_\mu \varphi_{\alpha\beta} = 0, \quad (4.80)$$

i. e. $\varphi_{\alpha\beta}$ are the functions only of the coordinates on the quotient space M/G .

The further development of a theory can be performed only due to additional assumptions about the structure of the matrix B_N^M , that cannot be described in terms of the basic principles, having made a base of our quantization scheme.

Let B_N^M has the simplest form $B_N^M = \delta_N^M$. In this case, as it follows from obtained above, the quantum mechanics is described by

$$[x^M, p_N] = i\hbar\delta_N^M, \quad [x^M, \dot{x}^N] = i\hbar\eta^{MN}. \quad (4.81)$$

The relations (4.81) can be divided into two parts. The first one corresponds to quantum mechanics on the orbit,

$$[x^\mu, p_\nu] = i\hbar\delta_\nu^\mu, \quad [x^\mu, x^\nu] = 0, \quad [p_\mu, p_\nu] = 0, \quad (4.82)$$

While the second one describes the quantum mechanics on the quotient space M/G :

$$[x^\alpha, x^\beta] = 0, \quad [x^\alpha, \pi_\beta] = i\hbar\delta_\beta^\alpha, \quad [\pi_\alpha, \pi_\beta] = i\hbar(p_\mu \circ F^\mu_{\alpha\beta} + \varphi_{\alpha\beta}). \quad (4.83)$$

The remaining commutation relations, containing in (4.81) have the form

$$\begin{aligned} [x^\alpha, x^\mu] &= 0, & [\pi_\alpha, p_\mu] &= -i\hbar\partial_\mu A_\alpha^\nu \circ p_\nu, \\ [x^\mu, \pi_\alpha] &= -i\hbar A_\alpha^\mu, & [x^\alpha, p_\mu] &= 0. \end{aligned} \quad (4.84)$$

It is essential, that the features of quantum mechanics on M/G depend on the tensor $\varphi_{\alpha\beta}$, which cannot be determined in our quantization scheme. This object can be viewed as the strength tensor of some abelian gauge field. To show this, we have to take into account that the commutator (4.74) can be rewritten in the simpler form due to the restriction $B_N^M = \delta_N^M$, namely

$$[p_\alpha, p_\beta] = i\hbar\varphi_{\alpha\beta}(x^\gamma). \quad (4.85)$$

Using (4.85) we find from the Jacobi identities for the operators $p_\alpha, p_\beta, p_\gamma$ that $\varphi_{\alpha\beta}$ obeys the relation

$$\partial_\alpha\varphi_{\beta\gamma} + \partial_\beta\varphi_{\gamma\alpha} + \partial_\gamma\varphi_{\alpha\beta} = 0. \quad (4.86)$$

Then, the evident solution of (4.86) is

$$\varphi_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha, \quad (4.87)$$

where A_α is some unknown abelian gauge field.

5 Quantum Lagrangian on M

Following [2] we introduce the following operator as the Lagrangian of a point particle with the unit mass

$$L(x, \dot{x}) := \frac{1}{2}(\dot{x}, \dot{x}) = \frac{1}{2}(p, p). \quad (5.88)$$

Here $(\cdot, \cdot)$ means the scalar product on M that is invariant under a general coordinate transformation $x \rightarrow \overline{x}(x)$. Its constructing has been detally discussed in [2]. In the present paper we modify this definition for the case

$$[x^M, p_N] = i\hbar B_N^M(x), \quad (5.89)$$

where $B_N^M \neq \delta_N^M$. One can simply show that B_N^M is the tensor field on M (and, consequently, $b_N^M = B_N^M - \delta_N^M$ is the tensor field too).

Under a general coordinate transformation $x \rightarrow \bar{x}(x)$ on M the momentum operator transforms as

$$p_M \rightarrow \bar{a}_M^N \circ p_N = \bar{a}_M^N p_N + \frac{1}{2}[p_N, \bar{a}_M^N] = \bar{a}_M^N p_N - \frac{i\hbar}{2}(\partial_N \bar{a}_M^N + b_M^P \partial_P \bar{a}_M^N), \quad (5.90)$$

$$p_M \rightarrow \bar{a}_M^N \circ p_N = p_N \bar{a}_M^N - \frac{1}{2}[p_N, \bar{a}_M^N] = p_N \bar{a}_M^N + \frac{i\hbar}{2}(\partial_N \bar{a}_M^N + b_M^P \partial_P \bar{a}_M^N), \quad (5.91)$$

Taking into account the transformation law of the Christoffel symbols Γ_{MN}^P , constructed with the metric η_{MN} , we can write down the derivatives of the transformation matrices as

$$\partial_N \bar{a}_M^N = \bar{\Gamma}_M + \bar{a}_M^N \Gamma_N, \quad \Gamma_N = \Gamma_{MN}^M, \quad (5.92)$$

$$\partial_N \bar{a}_M^P = \bar{\Gamma}_{LM}^S \bar{a}_N^L \bar{a}_S^P - \bar{a}_M^S \Gamma_{NS}^P. \quad (5.93)$$

Let us introduce the notation

$$W_M = \Gamma_{MN}^L b_L^N. \quad (5.94)$$

This object transforms under a general coordinate transformation on M as

$$\bar{W}_M = \bar{a}_M^N W_N + b_L^N \partial_N \bar{a}_M^L. \quad (5.95)$$

Hence, taking into account the transformation laws of p_M , Γ_M and W_M we can define the following “left” and “right” parts of the momentum operator

$$\begin{aligned} \pi_M &= p_M - \frac{i\hbar}{2}(\Gamma_M + W_M), \\ \pi_M^\dagger &= p_M + \frac{i\hbar}{2}(\Gamma_M + W_M), \end{aligned} \quad (5.96)$$

that transform as

$$\pi_M^\dagger \rightarrow \bar{\pi}_M^\dagger = \bar{a}_M^N \pi_N^\dagger, \quad \pi_M \rightarrow \bar{\pi}_M = \pi_N \bar{a}_M^N.$$

Following [2] we define the scalar norm of the momentum operator p_M on M in terms of the operators $\pi_M^\dagger$ and π_M as

$$(p, p) = \pi_M \eta^{MN} \pi_N^\dagger. \quad (5.97)$$

Using the obtained above, the Lagrangian can be written in the form

$$L = \frac{1}{2}(\pi_\alpha - \pi_\mu A_\alpha^\mu) g^{\alpha\beta} (\pi_\beta^\dagger - A_\beta^\mu \pi_\mu^\dagger) + \frac{1}{2} \pi_\mu g^{\mu\nu} \pi_\nu^\dagger. \quad (5.98)$$

In the simplest case $B_N^M = \delta_N^M$ we can write it as

$$L = \frac{1}{2}(p_M - \frac{i\hbar}{2}\Gamma_M) \eta^{MN} (p_M + \frac{i\hbar}{2}\Gamma_M) = \frac{1}{2} p_M \eta^{MN} p_N + \frac{\hbar^2}{4} \left(\partial_M \Gamma^M + \frac{1}{2} \Gamma_M \Gamma^M \right), \quad (5.99)$$

where

$$\Gamma_M = \Gamma_{MN}^N, \quad \Gamma^M = \eta^{MN} \Gamma_N. \quad (5.100)$$

The further analyzes will be performed in the restriction $B_N^M = \delta_N^M$. Define the metric on the group manifold G :

$$\eta_{ij} = g_{\mu\nu} v_i^\mu v_j^\nu, \quad \partial_M \eta_{ij} = 0. \quad (5.101)$$

The norm of the momentum operator on G

$$\mathcal{P}|_G = \{v_i^\mu \circ p_\mu : i = \overline{m=1, p}\}$$

is defined as

$$(\mathcal{P}|_G, \mathcal{P}|_G) = p_i \eta^{ij} p_j. \quad (5.102)$$

From the Killing equations in the form $\nabla_M v^M = 0$ it follows that

$$\partial_M v_A^M \equiv \partial_\mu v_A^\mu = -\Gamma_\mu v_A^\mu. \quad (5.103)$$

In the same time we conclude from (2.10) that

$$\Gamma_\mu = -e_\mu^i \partial_\nu v_i^\nu = v_i^\nu \partial_\nu e_\mu^i = \Lambda^\nu{}_{\nu\mu}. \quad (5.104)$$

Performing a direct calculation, one can obtain that

$$\Gamma_\alpha = \Gamma_{N\alpha}^M = \frac{1}{2} g^{\beta\gamma} \partial_\alpha g_{\beta\gamma} + \frac{1}{2} g^{\mu\nu} \partial_\alpha g_{\mu\nu}, \quad (5.105)$$

Finally, due to (5.104), (5.105) the definition (5.102) can be written as

$$(\mathcal{P}|_G, \mathcal{P}|_G) = \left(p_\mu - \frac{i\hbar}{2} \Gamma_\mu \right) g^{\mu\nu} \left(p_\mu + \frac{i\hbar}{2} \Gamma_\mu \right), \quad (5.106)$$

i. e. the norm of a group momentum coincides with the double Lagrangian for a free particle on the orbit of G on M .

Using (5.96) we can transform the Lagrangian into another form

$$L = \frac{1}{2} p_j g^{ij} p_j + \frac{1}{2} \left(p_\alpha - p_\mu A_\alpha^\mu - \frac{i\hbar}{2} \hat{\Gamma}_\alpha \right) g^{\alpha\beta} \left(p_\alpha - A_\alpha^\mu p_\mu + \frac{i\hbar}{2} \hat{\Gamma}_\alpha \right), \quad (5.107)$$

where

$$\hat{\Gamma}_\alpha = \Gamma_\alpha - A_\alpha^\mu \Gamma_\mu.$$

So we can rewrite (5.106) in terms of $\pi_\alpha = p_\alpha - A_\alpha^\mu \circ p_\mu$ as

$$L = \frac{1}{2} p_i \eta^{ij} p_j + \frac{1}{2} \left(\pi_\alpha - \frac{i\hbar}{2} \Omega_\alpha \right) g^{\alpha\beta} \left(\pi_\alpha + \frac{i\hbar}{2} \Omega_\alpha \right), \quad (5.108)$$

where

$$\Omega_\alpha = \hat{\Gamma}_\alpha - \partial_\mu A_\alpha^\mu.$$

In order to simplify (5.108) we have to present the object Ω_α in the explicit form. Using the structure equation we obtain

$$\partial_\mu A_\alpha^\mu = -A_\alpha^\nu \Lambda_{\nu\mu}^\mu + \Lambda_{\alpha\mu}^\mu, \quad (5.109)$$

and, consequently,

$$\Omega_\alpha = \Gamma_\alpha + A_\alpha^\mu \Lambda_{[\mu\nu]}^\nu - \Lambda_{\alpha\mu}^\mu = \Gamma_\alpha + v_i^\mu \partial_\alpha e_\mu^i + C^i_{ij} A_\alpha^j, \quad (5.110)$$

where

$$A_\alpha^i = e_\mu^i A_\alpha^\mu.$$

Further, taking into account (2.18) and the equality

$$\det \{\eta_{AB}\} = \det \{g_{\alpha\beta}\} \cdot \det \{v_i^\mu\}, \quad (5.111)$$

we arrive to the final expression for Ω_α :

$$\Omega_\alpha = \gamma_\alpha + \frac{1}{2} \eta^{ij} \partial_\alpha \eta_{ij} + C^i_{ij} A_\alpha^j, \quad (5.112)$$

where $\gamma_\alpha = \gamma^\beta_{\alpha\beta}$, $\gamma^\alpha_{\beta\gamma}$ denotes the Christoffel symbol constructed with the metric $g_{\alpha\beta}$ of the quotient space.

The formulae (5.108) and (5.112) completely determine the final expression for the Lagrangian for freely moving particle on M in the restriction $B_N^M = \delta_N^M$. The first term in (5.108) corresponds to quantum theory on the orbits of the action of G on M , While the second one describes the theory on the quotient space M/G . It is essential to point out here that the degrees of freedom $\{x^\mu\}$ determine the quantum mechanics on M/G by means of the last terms in (5.108).

6 Equations of Motion for Dynamics on Riemannian Manifold

In this section we restrict the function B_N^M to be equal to δ_N^M . The way of determining the dynamical equations of motion describing the particle moving on the manifold with an intransitive group of isometries differs from one introduced in [2] only in details. From the condition $\delta L = 0$ of permissible variations we find that

$$v_A^M \circ \left(\dot{p}_M + \frac{1}{2} p_L \partial_M \eta^{LN} p_N + \partial_M U \right) = 0, \quad (6.113)$$

here

$$U = \frac{\hbar^2}{4} \left(\partial_M \Gamma^M + \frac{1}{2} \Gamma_M \Gamma^M \right) \quad (6.114)$$

where we have substituted the Lagrangian (5.99).

Using the condition $v_i^\mu e_\nu^i = 0$, where $\{e_\mu^i\}$ is the inverse of $\{v_i^\mu\}$ we obtain the equations of motion of the free particle for degrees of freedom connected with the orbits of the action of G on M in the following form

$$\dot{p}_\mu = -\frac{1}{2}p_M\partial_\mu\eta^{MN}p_N - \partial_\mu U. \quad (6.115)$$

This equation is equivalent to the conservation law $\dot{p}_i = 0$ for the group momentum p^i .

As to the other degrees of freedom associated with M/G , its dynamical equations in Euler-Lagrange form must be equivalent to ones in Heisenberg form

$$\dot{p}_M = \frac{1}{i\hbar}[p_M, H]. \quad (6.116)$$

The explicit expression of the Hamiltonian can be obtained by comparison (6.115) with (6.116) taking $M := \mu$. The Hamiltonian operator derived in such a way has the following form

$$H = \frac{1}{2}p_M\eta^{MN}p_N + U + u_q(x^\alpha), \quad (6.117)$$

where $u_q = u_q(x^\alpha)$ is some function that does not appear in (6.115) (because $\partial_\mu u_q \equiv 0$). The detail calculation allow us to put $u_q = 0$ (see the remark below).

Hence, we can transform (6.113) into the following form

$$\dot{p}_M = -\frac{1}{2}p_L\partial_M\eta^{LN}p_N - \partial_M U - \left(\eta^{LN} \circ \varphi_{MN}\right) \circ p_L, \quad (6.118)$$

where

$$\varphi_{MN} = \frac{1}{i\hbar}[p_M, p_N].$$

The restriction of M to μ reduces (6.118) to (6.115), as it must be.

To prove the fact that (6.118) is correctly defined, we consider the operator H as the generator of time shifts $t \rightarrow \bar{t} = t + \delta t(t)$, where $\delta t(t)$ are permissible time variations. The result of the time variation of L must be in agreement with the equations of motion in Euler-Lagrange form (6.118).

The time shift causes the following variations of coordinate and velocity operators

$$\begin{aligned} \delta x &= \bar{x}(\bar{t}) - x(t) = \dot{x}(t) \delta t, \\ \delta \dot{x} &= \frac{d}{dt}\delta x - \dot{x} \frac{d\delta t}{dt}. \end{aligned} \quad (6.119)$$

Using the commutation relations we obtain

$$[x^M, \delta x^N] = i\hbar\eta^{MN}\delta t, \quad [\dot{x}^M, \delta x^N] = [\dot{x}^M, \dot{x}^N]\delta t, \quad (6.120)$$

where

$$[\dot{x}^M, \dot{x}^N] = i\hbar\varphi^{MN} + i\hbar f_L^{MN} \circ \dot{x}^L, \quad (6.121)$$

$$\begin{aligned} \varphi^{MN} &= \eta^{ML}\eta^{NS}\varphi_{LS}, \\ f^{MN}_P &= \left(\eta^{MS}\eta^{NL} - \eta^{NS}\eta^{ML}\right) \partial_S\eta_{LP}. \end{aligned} \quad (6.122)$$

The variation of the Lagrangian caused only by the time shift appears in the variation of the action functional in the following combination

$$\delta W = \int_{t_2}^{t_1} \delta_t L dt, \quad \delta_t L = \delta L + L \frac{d\delta t}{dt}. \quad (6.123)$$

The expression $\delta_t L$ reduces to

$$\delta_t L = \frac{d(L\delta t)}{dt}. \quad (6.124)$$

only due to the commutation relations.

On the other hand, extracting the total time derivative we obtain

$$\delta L = \frac{d}{dt} (H\delta t) + \frac{dL}{dt} - \left(\dot{p}_M - \frac{1}{2} p_N \partial_M \eta^{NL} p_L - \partial_M U - p_N \circ \varphi^\mu_M \right) \circ \delta x^N, \quad (6.125)$$

here H means the Hamiltonian. Hence, we can write

$$H = L, \quad \frac{dH}{dt} = 0, \quad (6.126)$$

after comparing (6.124) with (6.125). Note that if we define the Hamiltonian with an additional term $u_q(x)$, we obtain after such a comparison that $u_q = 0$ (otherwise the equation (6.124) and (6.125) are not agreed). Therefore, we can see that the equations (6.118) are consistent with other conclusions.

The equations of motion for degrees of freedom representing the orbits of the action of G on M are introduced above (see (6.115)). To write down the other dynamical equations describing the motion on M/G it is sufficient to assume $M \rightarrow \alpha \in \overline{1, m}$. In terms of π_α they are found in the form

$$\begin{aligned} \dot{\pi}_\alpha = & -\frac{1}{2} \pi_\beta \partial_\alpha g^{\beta\gamma} \pi_\gamma + \frac{1}{2} \left(\pi_\beta g^{\beta\gamma} F^\mu_{\alpha\gamma} p_\mu + p_\mu g^{\beta\gamma} F^\mu_{\alpha\gamma} \pi_\beta \right) \\ & - \frac{1}{2} p_\mu (g^{\mu\sigma} \partial_\sigma A^\nu_\alpha + g^{\nu\sigma} \partial_\sigma A^\mu_\alpha) p_\nu + p_M \circ \varphi^M_\alpha - \hat{D}_\alpha U \\ & + \frac{\hbar^2}{4} \hat{D}_\gamma (\partial_\alpha g^{\beta\gamma} \partial_\mu A^\mu_\beta) - \frac{\hbar^2}{8} \partial_\alpha g^{\beta\gamma} (\partial_\mu A^\mu_\beta \partial_\nu A^\nu_\gamma) \\ & + \frac{\hbar^2}{4} \partial_\mu (F^\mu_{\alpha\beta} g^{\beta\gamma} \partial_\nu A^\nu_\gamma) + \frac{\hbar^2}{4} \partial_M (\eta^{MN} \partial_M \partial_\mu A^\mu_\alpha). \end{aligned} \quad (6.127)$$

The equations (6.115), (6.127) completely describe the motion of a particle on the manifold with an intransitive group of motions. It appears that dynamics is decomposed into two parts. The first one corresponds to the motion on the orbits of the action of G on M , the equations of motion have the form of conservation laws for the group momenta $\{p_i : i = \overline{m+1, p}\}$. The second one is related to the motion on M/G , where the dynamics of a particle is governed by Lorentz-type forces. The first one is caused by non-abelian gauge field A^μ_α with the strength $F^\mu_{\alpha\beta}$, that has a purely geometrical origin, while the second one is related to the abelian gauge field φ_α with the strength $\varphi_{\alpha\beta}$ and appears due to quantum-mechanical nature of a theory (in the classical limit $\hbar \rightarrow 0$ this object vanishes). As to terms in (6.127) with the factor $\hbar^2$, they describes rather the operator orderings in the first terms in the r. h. s. of (6.127) than the features of dynamics.

7 Hilbert Space of States

Following the scheme introduced in [2] we chose the set of eigenvectors of coordinate operators $\{x^M\}$ as the basis of the Hilbert space for a point particle on M , namely

$$\hat{x}^M |x'\rangle = x'^M |x'\rangle . \quad (7.128)$$

with the following normalization condition

$$\langle x' | x'' \rangle = \Delta(x' - x'') := \frac{1}{\sqrt{\eta(x)}} \delta(x' - x'') , \quad (7.129)$$

where $\eta(x) = \det(\eta_{MN}(x))$. The Δ -function has the following properties

$$f(x') \Delta(x' - x'') = f(x'') \Delta(x' - x'') , \quad (7.130)$$

$$f(x') \partial'_M \Delta(x' - x'') = -\partial'_M f(x') \Delta(x' - x'') , \quad (7.131)$$

$$\partial'_M \Delta(x' - x'') = \partial'_M \left(\frac{1}{\sqrt{\eta(x')}} \Delta(x' - x'') \right) = -\Gamma_M(x') \Delta(x' - x'') - \partial''_M \Delta(x' - x'') . \quad (7.132)$$

In terms of coordinate decomposition $x^M = \{x^\alpha, x^\mu\}$ the equation (7.132) takes the form

$$\left(\hat{D}'_\alpha + \hat{D}''_\alpha + \hat{\Gamma}_\alpha(x') \right) \Delta(x' - x'') = \partial'_\mu A^\mu_\alpha(x') \Delta(x' - x'') , \quad (7.133)$$

where

$$\hat{D}_\alpha = \partial_\alpha - A^\mu_\alpha \partial_\mu , \quad \hat{\Gamma}_\alpha = \Gamma_\alpha - A^\mu_\alpha \Gamma_\mu .$$

To construct the coordinate representation of the operators describing the quantum mechanics on M one have to calculate the matrix elements of coordinate and momentum operators x^M and p_M .

In order to simplify this task we use previously defined operators

$$\bar{p}_M = \{\bar{p}_\alpha, \bar{p}_\mu\} , \quad \bar{p}_\alpha := p_\alpha - \varphi_\alpha , \quad \bar{p}_\mu := p_\mu , \quad (7.134)$$

that obey the following commutation relations

$$[x^M, x^N] = 0 , \quad [x^M, \bar{p}_N] = i\hbar \delta_N^M , \quad [\bar{p}_M, \bar{p}_N] = 0 . \quad (7.135)$$

Using the results of [2] we can write

$$\begin{aligned} \langle x' | x^M | x'' \rangle &= i\hbar x'^M \langle x' | x'' \rangle , \\ \langle x' | \bar{p}_M | x'' \rangle &= -i\hbar \left(\partial'_M + \frac{1}{2} \Gamma_M(x') \right) \Delta(x' - x'') . \end{aligned} \quad (7.136)$$

Returning to the operators p_M , we find that

$$\begin{aligned}
\langle x' | p_\alpha | x'' \rangle &= -i\hbar \left(\partial'_\alpha + \frac{1}{2} \Gamma_\alpha(x') \right) \Delta(x' - x'') + \varphi_\alpha(x') \Delta(x' - x''), \\
\langle x' | p_\mu | x'' \rangle &= -i\hbar \left(\partial'_\mu + \frac{1}{2} \Gamma_\mu(x') \right) \Delta(x' - x''), \\
\langle x' | x^M | x'' \rangle &= x''^M \Delta(x' - x'').
\end{aligned} \tag{7.137}$$

Using these expressions we can write the matrix element of the operator π_α in the form

$$\langle x' | \pi_\alpha | x'' \rangle = -i\hbar \left(\hat{D}_\alpha + \frac{1}{2} (\hat{\Gamma}_\alpha - \partial_\mu A_\alpha^\mu) \right) \Delta(x' - x'') + \varphi_\alpha(x') \Delta(x' - x''). \tag{7.138}$$

The coordinate representation of the operator A can be constructed in terms of wave functions $\psi(x') = \langle x' | \psi \rangle$ representing the state $|\psi\rangle$ by the usual way:

$$(\hat{A}\psi)(x') := \int \langle x' | A | x'' \rangle \langle x'' | \psi \rangle dx''. \tag{7.139}$$

Therefore, following [2] we find the representations for the coordinate and momentum operators in the following form

$$\begin{aligned}
\hat{x}^M \psi(x') &= x'^M \psi(x'), \\
\hat{p}_\alpha \psi(x') &= -i\hbar \left(\partial'_\alpha + \frac{1}{2} \Gamma_\alpha(x') \right) \psi(x') + \varphi_\alpha(x') \psi(x'), \\
\hat{p}_\mu \psi(x') &= -i\hbar \left(\partial'_\mu + \frac{1}{2} \Gamma_\mu(x') \right) \psi(x').
\end{aligned} \tag{7.140}$$

The coordinate representation of the Hamiltonian can be obtained in terms of the operators

$$\pi_M = p_M - \frac{i\hbar}{2} \Gamma_M, \quad \pi_M^\dagger = p_M + \frac{i\hbar}{2} \Gamma_M, \tag{7.141}$$

namely,

$$\hat{H} = \frac{1}{2} \hat{\pi}_M \hat{\eta}^{MN} \hat{\pi}_N^\dagger. \tag{7.142}$$

Using the decomposition of the coordinates of M we can write

$$\pi_\alpha = -i\hbar (\partial_\alpha + \Gamma_\alpha) + \varphi_\alpha, \quad \pi_\mu = -i\hbar (\partial_\mu + \Gamma_\mu), \tag{7.143}$$

$$\pi_\alpha^\dagger = -i\hbar \partial_\alpha + \varphi_\alpha, \quad \pi_\mu^\dagger = -i\hbar \partial_\mu. \tag{7.144}$$

Hence, the formal part of the formulation of the quantum mechanics on the Riemannian manifold with an intransitive group of isometries is solved. Nevertheless, the other one, that contains the meaning and interpretation of these results, requires the special discussion.

8 Discussion

We have examined in the present paper and in [2], [3] the generalization of Schwinger's action principle for the case of Riemannian manifolds with a group structure, determined by isometric transformations.

The obtained results show fundamental unity between geometrical properties of a manifold M and the algebra of quantum-mechanical operators, describing a point particle on M . The key role in the formulation of quantum mechanics is played by geometrical aspects of a symmetry, expressed in terms of Killing vectors. These vectors also determine the integrals of motion on M and their algebraic properties.

It appears that a general quantization problem is conditionally reduced into three cases. The first one we have considered in [2] is related to the class of manifolds with a simply transitive group of isometries. Here the dimension of this group (which coincides with the number of Killing vectors) is exactly equals to the dimension of the manifold. The quantum mechanics receives its traditional form, investigated in [4], [5].

The second case contains the investigation of homogeneous manifolds with non-simply transitive group of isometries, where the dimension of a group exceeds the dimension of a manifold. There appears a non-abelian gauge structure, that is similar to one in Kaluza-Klein theories. This structure is related to the isotropy subgroup of G and plays the role of gauge group. This case has been analyzed in [3]. The obtained results are in accordance with [6], where the method of study somewhat differs from ours.

The remaining third case, analyzed in the present paper, is related to Riemannian manifolds with the intransitive group of isometries G . In this case the dimension of the manifold M is bigger than the dimension of the group G . From the geometrical point of view M can be considered as a total space of the principal fiber bundle with the base space M/G and the structure group G . In terms of this construction M can be covered by the set of submanifolds (the orbits of G -action) and G acts simply transitively in each of them. Quantum mechanics on such a submanifold is developed in [2]. The other degrees of freedom, related to the quotient space M/G have not uniquely determined dynamics. In order to determine it unequivocally we have to assume something new and external in respect to Schwinger's scheme. There are two types of arbitrary factors in the formulation of quantum mechanics on M . Some of them have geometrical nature (the functions b_N^M), while the others are quantum mechanical objects (the abelian gauge field φ_α that vanishes when $\hbar \rightarrow 0$). It is important to point out here, that the last structure can not be obtained in the framework of canonical quantization procedure, it is caused by geometrical properties of the manifold and algebraic relation between geometrical objects.

The formal results of the present paper bring up the more complicated question about its interpretation. We suppose that it can be done in the following manner. There are some degrees of freedom, describing a quantum system, which have partially undetermined properties (as it can be treated in the traditional meaning). The geometry of space is determined by energy-momentum of the matter distribution. It is substantiated to search the connection between geometrical structures corresponding to undetermined dynamics and forbidden spatial zones for the system with discrete energy spectrum (in the manner of "layers" between the Bohr orbits in the hydrogen atom). Nevertheless, the question about the interpretation of quantum theory on a general Riemannian manifold remains an open one and requires more detailed discussion and further development.

As the logical continuation of our series of papers we thought the generalization of Schwinger's quantization approach on the case of a supermanifold, that we are going to present in the forthcoming paper.

References

- [1] L. Eisenhart, *Continous Groups of Transformations*, Princeton, 1933
- [2] N. Chepilko, A. Romanenko, Submitted to Europ. Phys. Journ.
- [3] N. Chepilko, A. Romanenko, Submitted to Europ. Phys. Journ.
- [4] R. Sugano, *Prog.Theor.Phys.* **46**, 297 (1971);
T. Kimura, *ibid.* **46**, 126 (1971);
R. Sugano and T. Kimura, *ibid.* **47**, 1004 (1972);
T. Ohtani and R. Sugano, *ibid.* **47**, 1704 (1972); *ibid.* **50**, 1705 (1973)
- [5] B. S. DeWitt, *Rev. Mod. Phys.* **29**, 377 (1957).
- [6] D. McMullan and I. Tsutsui, *Ann. Phys.* **237**, 269 (1995);
Y. Ohnuki and S. Kitakado *J. Math. Phys.* **34**, 2827 (1993).